



सत्यमेव जयते

**Ministry of Women and Child Development
Government of India**

Rapid Survey on Children 2013-2014

INDIA Fact Sheet

The results of **Rapid Survey on Children (RSOC)** are intended to strengthen the data system on children and women, based on a nationwide household cum facility based survey in 28 states and Delhi. This survey provides level of selected indicators at national and state level. The indicators mainly focus on the wellbeing of children below 6 years and their mothers covering aspects of child development, maternal care, school /college attendance among persons aged 5-24 years. Early childhood care and the enabling environment, like access to drinking water and use of toilet facilities are also covered in the survey. The survey maps the ICDS programme in terms of its infrastructural facilities, awareness and utilization of the six services by the target groups.

Respondents: Head of the selected household, all ever married women (EMW) aged 15-49 years; currently married women aged 15-49 pregnant at the time of survey in the selected households and Anganwadi workers (AWW) in the selected AWC.

Selection of Primary Sampling Unit (PSU), Household and AWC: Selection of PSUs was similar to the National Family Health Survey 2005-06. PSUs were selected using Probability Proportional to Size

(PPS) with replacement and from each PSU, selection of 26 households was done by circular systematic random sampling in a manner that 20 households from households having at least one child below 6 years of age and 6 households from other households. All AWC in the selected PSU / Segmented PSUs were selected in the sample. Overall 2655 rural PSUs and 1497 urban PSUs were covered.

RSOC covered 105,483 households and 5630 Anganwadi centres (AWC) leading to more than 210,000 interviews. For compiling nutritional status, measurement of height and weight of over 90,000 children aged 0-4 and over 28,000 adolescent girls aged 10-18 have been collected across all states. The anthropometric equipment used for the measurement are of international specifications. More than 167 exclusive health investigators were trained and engaged for taking anthropometric measurements in this survey.

RSOC fieldwork was completed in 23 weeks (3rd week of November 2013 to 2nd week of May 2014), by 188 listing teams and 167 survey teams. The survey was conducted on CAPI mode.





Key Indicators		Residence			Gender		Social category				Wealth index	
		Total	Rural	Urban	Male	Female	SC	ST	OBC	Other	WI- Lowest	WI- Highest
A	Women who had live-birth in 35 months prior to survey by specific maternal health care (Percentage)											
1	Prenatal Care¹											
1.1	Registered pregnancy	84.1	82.7	87.4			82.0	86.1	82.4	87.1	75.9	89.1
1.2	Received Mother and Child Protection (MCP) card at the time of pregnancy registration	73.0	72.0	75.4			71.8	71.2	70.8	77.6	62.0	78.2
1.3	Received at least one ante-natal check-up (ANC)	85.2	83.0	90.3			83.8	81.8	84.2	88.8	74.7	94.0
1.4	Received first ANC in first trimester	61.8	58.5	69.7			59.6	54.7	62.8	65.1	47.0	77.0
1.5	Received three or more ANCs	63.4	58.5	74.9			58.9	56.4	60.9	72.9	43.7	80.1
1.6	Received four or more ANCs	45.4	39.6	58.9			41.8	37.0	43.3	54.4	24.5	64.4
1.7	Received two or more Tetanus Toxoid (TT) injections	89.8	88.8	92.2			89.2	86.1	89.4	92.3	84.4	94.8
1.8	Received/purchased 100 or more IFA tablets /syrup during pregnancy	31.2	29.1	36.1			29.3	30.5	30.3	34.3	20.7	41.8
1.9	Consumed 100 or more IFA tablets /syrup during pregnancy	23.6	21.4	28.8			21.6	20.6	23.6	26.6	13.8	35.0
1.10	Received full ANC ²	19.7	17.3	25.2			18.0	15.0	19.6	23.2	9.5	31.3
1.11	Visited at least once during pregnancy ³ by											
	ANM	13.9	14.9	11.4			14.3	13.8	14.3	13.1	10.5	11.1
	AWW	34.8	38.7	25.7			36.1	42.5	33.6	32.5	33.5	27.2
	ASHA	31.6	39.4	13.3			33.3	37.4	29.4	30.6	36.8	19.7
1.12	Received ANC at											
	Anganwadi centre	22.4	27.4	10.6			24.3	36.5	23.0	14.4	32.4	12.3
	Government health facility	43.3	42.7	44.6			48.8	43.1	38.0	46.1	37.8	36.1
	Private health facility	38.1	33.1	49.7			30.3	22.8	41.1	45.7	19.3	63.9
2	Natal Care											
2.1	Institutional delivery	78.7	74.6	88.5			76.0	70.1	79.2	84.2	60.8	93.0
2.2	Delivered by skilled health provider	81.1	77.2	90.2			78.8	72.7	81.6	86.1	64.1	94.5
2.3	Stay at health facility after delivery											
	Less than 12 hours	20.6	25.1	11.7			25.4	17.0	22.0	16.8	32.9	12.6

¹ It is based on births taken place during the last three years

² Full ANC- receipt of 3 + ANC, atleast 1 dose of TT and consumption of 100 IFA tablets/ syrup

³ Out of those that have AWC in their area



Key Indicators		Residence			Gender		Social category				Wealth index	
		Total	Rural	Urban	Male	Female	SC	ST	OBC	Other	WI- Lowest	WI- Highest
	48 hours or more	57.8	51.5	70.2			54.1	55.1	58.8	60.1	39.1	70.1
3	Post Natal Check-up (PNC)											
3.1	Received PNC within 48 hours of discharge/delivery											
	All	39.3	34.2	51.1			35.3	36.6	43.3	38.0	23.0	48.8
	Home delivery (after birth)	13.0	12.4	15.9			12.8	12.1	15.4	9.5	9.9	23.6
	Institutional delivery (after discharge)	46.3	41.6	55.7			42.4	47.0	50.6	43.3	31.5	50.8
3.2	New born who received first check up within 24 hours of birth/discharge											
	All	33.6	31.0	39.8			31.2	34.0	35.8	32.6	22.9	37.6
	Home delivery (after birth)	13.0	12.6	15.0			13.2	11.6	15.4	9.6	10.0	22.9
	Institutional birth (after discharge)	39.2	37.2	43.1			36.9	43.5	41.1	36.9	31.2	38.7
3.3	Visited by primary health worker (AWW/ ANM/ ASHA) at home within one week of delivery/ discharge from health Institution	51.0	49.4	54.7			49.0	53.5	53.5	48.3	40.9	53.2
3.4	Availed benefit from national programme for safe motherhood ⁴											
	Janani Suraksha Yojana (JSY)	47.7	51.8	36.6			54.5	54.7	47.7	39.4	54.3	31.9
	Janani Shishu Suraksha Karyakrama (JSSK)	14.0	14.2	13.6			15.2	16.1	12.8	14.2	11.6	11.0
	Both JSY and JSSK	7.8	8.3	6.5			9.6	8.4	7.3	7.0	7.1	5.8
B	Birth Weight (Percentage of children aged 0 –35 months)											
1	Weighed with 24 hours of birth	68.7	64.1	79.7			65.3	63.7	67.4	75.5	49.9	83.0
2	With birth weight less than 2500 grams (out of those weighed)	18.6	18.7	18.4			19.6	21.6	18.0	17.6	22.3	15.6
C	Immunization (Percentage of children aged 12 –23 months)											
1	Percentage of children aged 12-23 months											
1.1	Having MCP/ immunization card	84.3	82.8	87.8	84.4	84.2	83.5	79.3	82.9	89.0	75.6	90.9
1.2	Fully immunized ⁵	65.3	62.4	72.0	65.0	65.7	61.6	55.7	65.4	71.6	50.6	80.0
1.3	Received DPT 3 injection	74.8	72.2	80.7	74.3	75.3	72.0	65.0	74.9	80.5	62.4	87.2
1.4	Received first dose of Measles vaccine	78.9	77.0	83.1	79.4	78.3	76.7	75.4	77.5	83.8	66.5	88.5
1.5	Received no vaccination	6.6	7.6	4.5	6.6	6.7	8.8	7.4	7.1	4.1	12.6	2.0
2	Drop out in vaccination											

⁴ Out of mothers who are aware JSY / JSSK schemes

⁵ Full Immunization- BCG, Measles, 3 doses each of DPT and Polio vaccine (excluding polio vaccine given at birth)



Key Indicators		Residence			Gender		Social category				Wealth index	
		Total	Rural	Urban	Male	Female	SC	ST	OBC	Other	WI- Lowest	WI- Highest
	DPT 2 to DPT 3	11.1	12.6	8.0	11.7	10.5	12.4	17.3	10.6	8.7	16.9	6.0
	OPV 2 to OPV 3	11.9	12.8	9.9	12.4	11.4	13.6	16.9	11.7	9.1	17.4	6.7
3	Place of the vaccination											
	Anganwadi centre	49.3	59.7	26.3			55.1	63.1	52.4	36.2	69.7	25.8
	Government health facility	35.3	29.9	47.2			36.3	27.7	29.6	44.7	23.1	40.5
	Private health facility	11.7	5.8	24.6			5.5	6.3	13.3	16.0	2.1	30.7
D	IYCF (Infant and Young Child Feeding Practices) (Percentage)⁶											
1	Children aged 0-23 months breastfed immediately/ within an hour of birth	44.6	44.2	45.6	43.8	45.4	43.1	54.7	42.3	44.8	42.5	42.3
2	Children aged 0-5 months who were exclusively breastfed ⁷	64.9	65.1	64.2	65.0	64.8	67.1	64.3	64.0	64.5	70.0	61.7
3	Children aged 6-8 months who were fed complementary foods	50.5	47.1	58.1	52.9	47.8	45.2	45.5	50.8	55.6	41.2	62.1
4	Breastfed children (6 -23 months)											
4.1	Fed a minimum number of times ⁸	36.3	35.7	37.7	36.8	35.8	35.8	37.8	36.8	34.7	35.9	40.1
4.2	Had a minimum dietary diversity ⁹	19.9	18.6	23.0	20.7	19.0	17.9	16.6	19.1	24.1	14.2	26.5
E	Nutritional Status of Children (Percentage)¹⁰											
1	Percentage of children aged 0-59 months											
1.1	Stunted (Height for age below -2SD)	38.7	41.6	32.0	39.5	37.8	42.4	42.3	38.9	33.9	50.7	26.7
1.2	Severely stunted (Height for age below -3SD)	17.3	19.1	13.2	17.6	16.9	19.3	19.5	17.8	14.2	25.8	10.7
1.3	Wasted (Weight for height below -2SD)	15.1	15.1	15.0	15.6	14.5	15.5	18.7	14.8	13.6	17.0	13.0
1.4	Severely wasted (Weight for height below -3SD)	4.6	4.5	4.8	4.8	4.4	4.9	5.3	4.4	4.4	5.0	4.4
1.5	Underweight (Weight for age below -2SD)	29.4	31.6	24.3	30.0	28.7	32.7	36.7	29.3	23.6	42.1	18.6
1.6	Severely underweight (Weight for age below -3SD)	9.4	10.6	6.9	10.0	8.9	10.8	13.0	9.3	7.0	16.3	5.1
F	Micronutrient and Deworming (Percentage)											

⁶ Based on standard set of questions as given in 'Indicators for Assessing IYCF practices part 2 Measurement' 2010 by USAID, WHO, UNICEF etc.

⁷ Exclusive breastfeeding- children who are fed breast milk exclusively for first six months (excluding pre-lacteals given at birth) and including ORS/medicine during sickness;

⁸ At least twice a day for breastfed infants 6-8 months old; at least three times a day for breastfed children 9-23 months old;

⁹ Minimum Dietary Diversity refers to four or more food groups to children age 6-23 months.

¹⁰ Based on WHO Growth Standards



Key Indicators		Residence			Gender		Social category				Wealth index	
		Total	Rural	Urban	Male	Female	SC	ST	OBC	Other	WI- Lowest	WI- Highest
1	Percentage of children aged 6-59 months received in six months prior to survey											
1.1	Vitamin A dose	45.2	43.8	48.3			43.4	46.0	45.5	46.4	37.5	48.8
1.2	Iron and Folic Acid supplement	13.4	12.7	15.2			12.3	15.4	13.7	13.5	8.7	15.1
1.3	Deworming medication	27.6	26.6	30.0			24.4	28.2	25.5	32.8	23.0	31.2
G	Morbidity - Percentage of children aged 0-59 months (Percentage)											
1	Diarrhoea											
1.1	Had diarrhoea in 15 days prior to survey	6.5	6.5	6.3	6.5	6.4	6.7	6.3	7.0	5.7	6.8	5.8
1.2	For whom advice or treatment was sought	77.9	78.5	76.4	79.8	75.7	78.9	73.0	77.3	79.8	77.2	77.6
1.3	Children with diarrhoea given											
	Oral Re-hydration Solution (ORS) including Home Available Fluids (HAF)	54.4	51.7	60.9	56.3	52.4	56.2	51.4	52.9	56.8	48.1	61.8
	ORS and Zinc	12.6	11.2	16.1	12.4	12.8	12.8	13.2	12.3	13.0	9.2	14.8
2	Fever											
2.1	Had fever in 15 days prior to survey	13.6	13.8	13.2	13.8	13.5	14.4	11.6	13.8	13.5	14.8	12.4
2.2	Given anti-malarial drug during fever	18.4	19.2	16.4	19.2	17.5	18.9	22.6	19.8	15.1	19.0	17.5
3	Prevalence of symptoms of ARI											
3.1	Had symptoms of ARI in 15 days prior to survey	8.6	9.1	7.6	8.9	8.3	9.0	7.9	8.8	8.3	10.6	7.0
3.2	For whom advice or treatment was sought	76.9	75.8	79.9	78.1	75.4	78.0	66.4	78.1	78.1	73.9	83.2
H	Pre-School Education and Literacy (Percentage)											
1	Percent of children aged 3-6 years currently attending pre-school education (PSE) at¹¹											
	Anganwadi centre	37.9	45.1	21.6	36.6	39.3	41.6	50.9	34.8	33.9	51.4	15.4
	Privately run institution (Play school, nursery etc.)	30.7	22.0	50.3	31.7	29.6	25.0	17.3	32.1	39.1	8.6	61.5
	Not attending	26.9	28.0	24.5	27.3	26.6	29.1	27.3	28.3	22.9	34.7	19.9
2	Pre-School Education (PSE)											
	Percentage of children aged 36-71 months attended PSE in AWC for 16 or more days in the month prior to survey (among	58.1	58.6	55.5	57.2	58.9	57.2	58.0	58.4	58.9	56.6	56.6

¹¹ Out of those households which are covered by an AWC



Key Indicators		Residence			Gender		Social category				Wealth index	
		Total	Rural	Urban	Male	Female	SC	ST	OBC	Other	WI- Lowest	WI- Highest
	children attending PSE at AWC)											
3	Literacy Rate 5 Years & above	71.9	67.0	82.2	79.0	64.6	65.9	61.9	72.2	79.3	52.1	88.6
4	Literacy Rate 6 Year & above											
5	Literacy Rate 7 Year & above	73.4	68.5	83.7	81.0	65.8	67.4	63.2	73.9	80.7	53.5	90.0
6	Youth Literacy rate (Aged 15-24)	88.7	87.0	92.5	91.9	85.6	86.2	81.5	89.1	92.8	74.3	97.5
I	Birth Registration and Marriage (percentage)											
1	Children aged below 5 Years whose birth is registered											
1.1	Birth registered	72.0	67.1	83.2	71.3	72.7	67.5	70.5	68.4	80.8	56.8	86.2
1.2	Have birth certificate	37.2	32.6	47.5	36.3	38.1	36.5	32.2	34.2	43.7	21.7	48.9
2	Mean age at marriage											
	Female	21.1	20.5	22.2			20.5	21.4	20.9	21.6	20.2	22.8
	Male	25.0	24.3	26.3			24.5	24.7	24.8	25.9	23.5	26.9
3	Women aged 20-24 married before age 18	30.3	33.5	22.4			34.9	31.0	30.5	25.7	44.1	15.4
4	Men aged 25-29 married before age 21	30.2	34.3	21.1			33.8	37.1	30.3	24.1	47.3	17.6
J	Adolescent Girls (10-19 years) (Percentage)											
1	Adolescent girls ever married											
	Age 10-19	6.4	7.2	4.5			7.5	6.4	6.0	5.8	6.7	3.2
1.1	Aged 10-14	0.3	0.4	0.1			0.5	0.6	0.3	0.1	0.4	0.2
1.2	Aged 15-19	12.8	14.4	8.9			15.1	12.7	12.0	11.5	15.3	5.9
2	Girls aged 15 -18 whose Body Mass Index (BMI) was (Height/Weight)¹²											
2.1	Less than 18.5 (low weight)	44.7	44.9	44.2			46.7	49.2	44.7	40.7	45.9	41.1
2.2	More than 25 (over weight)	4.5	3.1	7.9			3.2	4.4	4.3	5.9	2.9	6.0
K	Awareness and use of ICDS services (Percentage)											
1	Mothers of children aged 0-35 months aware of services at AWC											
	Supplementary food	85.1	87.2	79.3			86.5	89.2	84.0	83.6	86.0	79.1
	Immunization	57.4	59.2	52.5			56.0	62.8	61.8	50.9	55.3	58.3
	Health check-up	29.4	28.8	31.0			27.6	29.8	28.7	32.0	21.0	33.3
	Referral service	12.2	11.8	13.3			11.6	11.9	13.5	11.2	7.1	14.8
	Nutrition and health education	17.2	16.7	18.4			15.6	17.6	17.9	17.6	10.5	20.7

¹² BMI in kg/m²



Key Indicators		Residence			Gender		Social category				Wealth index	
		Total	Rural	Urban	Male	Female	SC	ST	OBC	Other	WI- Lowest	WI- Highest
	All Services ¹³	9.3	9.0	10.3			8.6	9.4	10.2	8.8	4.4	11.6
2	Mothers of children aged 36-71 months aware of services at AWC											
	Supplementary food	91.0	91.5	89.5			91.5	93.1	89.5	91.7	90.7	89.5
	Immunization	59.8	60.1	59.0			56.3	64.7	64.5	54.3	55.8	65.5
	Pre-School Education	49.9	49.3	51.6			46.3	50.1	49.4	54.0	42.1	56.3
	Health check-up	30.5	28.9	35.2			27.6	31.6	29.8	34.1	21.6	37.1
	Referral service	13.4	12.3	16.4			12.2	12.6	14.1	13.9	7.4	17.8
	Nutrition and health education	18.6	17.3	22.3			16.4	18.1	19.2	20.1	10.8	24.5
	All Services ¹⁴	9.7	8.8	12.2			8.4	9.1	10.1	10.6	4.2	13.7
3	Currently pregnant women aware of services at AWC											
	Supplementary food	87.7	88.3	85.8			88.4	91.0	86.2	87.6	86.4	87.3
	Immunization	59.6	59.9	58.6			59.1	62.5	62.7	55.4	53.5	62.5
	Health check-up	29.9	28.6	34.3			33.1	27.7	27.2	33.7	20.8	34.0
	Referral service	11.0	9.8	14.9			11.4	11.9	11.2	10.2	5.6	16.3
	Nutrition and health education	16.5	15.7	19.3			19.0	14.2	16.2	16.9	10.2	18.3
	All Services ¹⁵	8.3	7.2	11.9			9.5	9.0	7.8	8.0	2.6	10.8
4	Percentage availing various Anganwadi services											
4.1	Supplementary food											
	Children aged 6-35 months	49.2	53.8	36.7			53.0	62.8	45.1	45.8	57.0	32.0
	Children aged 36-71 months	44.2	48.7	31.1			46.6	57.9	39.1	43.0	53.7	24.0
	Pregnant women	40.7	44.0	29.8			43.9	55.6	36.7	36.7	43.7	27.1
	Lactating mothers	42.4	46.7	28.7			46.7	53.8	40.4	35.9	53.1	26.8
4.2	Percentage who received supplementary nutrition for 21 or more days in the month prior to survey ¹⁶											
	Children aged 6-35 months	21.3	20.5	24.3			20.8	24.8	23.7	16.7	16.8	28.0
	Children aged 36-71 months	31.0	30.9	31.7			31.6	33.1	29.8	31.3	29.9	33.9

¹³ All ICDS services as applicable for the mothers of children aged 0-35 months

¹⁴ All ICDS services as applicable for the mothers of children aged 36-71 months

¹⁵ All ICDS services as applicable for the currently pregnant women

¹⁶ Calendar month prior to survey refers to September, 2013



Key Indicators		Residence			Gender		Social category				Wealth index	
		Total	Rural	Urban	Male	Female	SC	ST	OBC	Other	WI- Lowest	WI- Highest
	Pregnant women	27.8	27.0	31.4			37.1	26.8	23.3	25.4	26.6	29.3
	Lactating mothers	20.3	20.5	18.9			22.3	26.0	19.9	15.8	16.5	26.4
L	Household characteristics (Percentage)											
1	Access to any improved source of drinking water	91.0	90.2	92.8			92.7	85.3	90.3	93.1	88.7	91.2
2	Using improved sanitation facility ¹⁷	41.8	29.9	66.0			29.8	22.2	43.3	55.4	4.5	84.5
3	Households practicing open defecation	45.5	61.6	12.8			58.1	68.9	45.7	28.0	91.1	2.3
4	Households using adequately iodized salt (>=15 ppm)	67.4	61.3	79.7			61.1	62.1	65.6	75.8	49.6	86.6

¹⁷ Using Improved sanitation means households using improved toilets as per JMP definition and this excludes those households, who are using improved toilets but sharing toilet facilities with other households



Key Indicators on Anganwadi Centre

Key Indicators		Total	Rural	Urban
CHARACTERISTICS				
A	General Information about Anganwadi worker (AWW) and AWC (Percentage)			
1	AWWs living in the AWC village/ward	74.1	75.5	60.9
2	AWW having 10 or more years of schooling	84.8	83.8	94.4
3	AWW below 45 years	78.4	78.9	74.7
4	Median age of AWW	38.0	37.0	39.0
5	AWCs serving to population more than the stipulated norm	56.9	54.5	78.0
6	AWW Received Flexi fund	43.9	43.5	48.2
B	Infrastructure Facilities at AWC (Percentage of AWC)			
1	AWC functioning in			
	Own ICDS building	40.5	42.8	19.6
	Rented building	21.3	17.1	60.9
	Out of those not functioning from ICDS or rented building			
	School building	49.8	50.9	26.4
	Panchayat Building		12.8	
	Anganwadi worker or Anganwadi helper's house	11.1	10.1	32.8
2	Various infrastructure facilities in AWCs			
2.1	Separate kitchen (among those centres that are cooking supplementary food in the AWC)	52.7	52.6	54.5
2.2	Cooking supplementary food in the same room where PSE sessions are conducted ¹⁸	15.9	15.4	21.9
2.3	Cooking supplementary food in open space	19.9	20.5	13.0
2.4	Barrier free access for physically challenged children	25.3	25.6	22.3
2.5	Having toilet facility	43.4	42.0	56.7
2.6	Access to drinking water			
	Within the premises of AWC	44.7	43.5	56.1
	Outside the premises of AWC but within 50 meters of AWC	62.9	61.9	74.2
2.7	Having electricity connection	32.4	28.6	68.0
3	Percentage of AWCs open for at least for 4 hours per day	93.1	92.8	96.2
4	AWCs functioning in rooms (other than kitchen space)			
	one room	65.6	64.5	75.0
	Two or more rooms	25.5	26.1	20.2
5	AWCs providing services			
	Supplementary nutrition	96.7	96.5	98.0

¹⁸ Out of those AWCs where Supplementary food is cooked in the AWC



Key Indicators		Total	Rural	Urban
	Pre-School Education	90.6	90.3	92.9
	Immunization	82.1	82.2	81.1
	Nutrition and health education	64.7	63.4	77.2
	Health check-up	60.8	60.1	66.9
	Referral	43.1	42.3	50.4
	Four or more services	71.5	70.6	80.2
	All six services	29.3	28.8	34.7
6	Percentage of AWCs functioned for 25 or more days in the calendar month prior to survey	62.6	63.1	57.4
C Supplementary Nutrition (Percentage)				
1	Coverage of Beneficiaries (Supplementary Food)			
	Children aged 6-35 months	78.3	80.2	76.4
	Children aged 36-71 months	73.6	74.4	67.0
	Pregnant women	83.1	83.4	82.8
	Lactating mothers	81.7	82.7	80.7
2	Beneficiaries received supplementary food for at least 21 days in the month ¹⁹ prior to survey			
	Children aged 6 -35 months	28.6	24.2	33.1
	Children aged 36-71 months	86.8	90.1	83.6
	Pregnant women	34.8	28.8	40.7
	Lactating mothers	33.8	28.7	38.9
D Growth Monitoring (Percentage of AWCs)				
1	AWCs having functional weighing scale			
	Baby weighing scale	72.4	72.4	72.1
	Adult weighing scale	51.6	51.3	53.8
2	Availability of WHO growth chart at AWC ²⁰	89.7	89.9	88.1
3	Children for whom Growth chart was prepared			
	0 -35 months	43.8	43.1	49.5
E Pre-School Education				
1	Average number of days spent on PSE sessions in the month prior to survey ²¹ (Days)	23.7	23.7	23.4

¹⁹ Month prior to survey refers to September, 2013

²⁰ Includes both- shown and not shown

²¹ Month prior to survey refers to September, 2013



Key Indicators		Total	Rural	Urban
2	Percentage of AWCs conducted PSE sessions in one calendar month prior to survey			
2.1	One or more hours a day	86.4	86.4	86.4
2.2	Sixteen or more days	95.2	95.3	94.4
3	Percentage of children aged 36-71 months attended PSE for 16 or more days in one calendar month prior to survey			
	Boys	55.9	57.1	46.8
	Girls	57.2	58.3	48.5
4	Average number of activities per day conducted in the week prior to survey	4.8	4.8	5.0
F	Convergence with health (Percentage of AWCs)			
1	AWCs that organized three Village Health and Nutrition Day (VHND) meetings during three calendar months prior to survey ²²	65.2	65.2	
2	AWCs reporting three visits by ANM during the three months prior to survey	83.6	83.6	
3	AWCs having Anganwadi level Management Committee (ALMC)	50.7	52.1	37.8
4	AWW reporting support from Panchayat/ward member	43.2	44.2	33.5
G	Training & Comprehensive Knowledge (Percentage of Anganwadi workers)			
1	Trainings received by AWWs			
1.1	Refresher training in two years prior to survey	42.5	42.5	41.8
	Special Training			
	IMNCI	50.5	50.2	52.8
	IYCF	55.1	54.8	58.7
	PSE	69.5	69.5	69.9
	Growth monitoring	59.6	59.3	62.2
	Early childhood care and development	49.1	48.6	53.6
	Joint ICDS-Health Training	44.4	43.9	48.3
	Revised ICDS MIS training	30.8	30.4	34.4
2	AWW having correct knowledge of			
2.1	Intake of food by Pregnant women	91.4	91.1	94.1
2.2	Normal Birth weight of a child	80.4	80.1	82.8
2.3	Initiation of breast feeding within one hour	96.0	96.1	95.3
2.4	Exclusive breast feeding for the first six months	92.9	92.7	94.9
2.5	Appropriate age of child for complementary food	63.8	63.6	65.8

²² Three Calendar month prior to survey refers to July, August, September, 2013



Population, sample coverage on key parameters and survey duration RSOC, 2013-14	
Population (Census 2011)	1,210,854,977
Sample coverage	
Number of primary sampling units (PSUs) covered	4,162
Households interviewed	105,483
Number of ever married women aged 15-49 years interviewed	111,636
Number of children 0-59 months for whom anthropometric measurements done	90,665
Number of Adolescent girls aged 10-18 years for whom anthropometric measurements done	28,522
Number of children aged 12-23 months	17,311
Month and year of field work	
From	11/2013
To	5/2014



Sampling errors of selected indicators, RSOC 20013-14

Key Indicators	Value (R)	Standard error (SE)	Design effect (DEFT)	Relative standard error (SE/R)	Confidence limits	
					R-2SE	R+2SE
Received at least one ante-natal check-up (ANC)	0.852	0.002	1.312	0.002	0.848	0.856
Received three or more ANCs	0.634	0.003	1.347	0.004	0.628	0.639
Consumed 100 or more IFA tablets /syrup during pregnancy	0.236	0.002	1.355	0.011	0.231	0.241
Institutional delivery	0.787	0.002	1.331	0.003	0.783	0.792
Delivered by skilled health provider	0.811	0.002	1.352	0.003	0.807	0.816
Fully immunized (children age 12-23 months)	0.653	0.005	1.395	0.008	0.643	0.663
Received BCG (children age 12-23 months)	0.911	0.003	1.455	0.003	0.905	0.917
Received DPT 3 injection (children age 12-23 months)	0.748	0.005	1.398	0.006	0.739	0.757
Received first dose of Measles vaccine (children age 12-23 months)	0.789	0.004	1.397	0.005	0.780	0.797
Children aged 0-5 months who were exclusively breastfed	0.649	0.007	1.417	0.011	0.635	0.663
Stunted (Height for age below -2SD) (children age 0-59 months)	0.387	0.002	1.460	0.006	0.382	0.392
Severely stunted (Height for age below -3SD) (children age 0-59 months)	0.173	0.002	1.448	0.011	0.170	0.177
Wasted (Weight for height below -2SD) (children age 0-59 months)	0.150	0.002	1.473	0.012	0.147	0.154
Severely wasted (Weight for height below -3SD) (children age 0-59 months)	0.046	0.001	1.474	0.023	0.044	0.048
Underweight (Weight for age below -2SD) (children age 0-59 months)	0.294	0.002	1.434	0.007	0.290	0.298
Severely underweight (Weight for age below -3SD) (children age 0-59 months)	0.094	0.001	1.454	0.015	0.092	0.097
Given vitamin A supplement in last 6 months (children age 6-59 months)	0.452	0.002	1.331	0.005	0.447	0.456
Had diarrhoea in 15 days prior to survey (children age 0-59 months)	0.065	0.001	1.464	0.018	0.062	0.067
Children with diarrhoea given ORS (children age 0-59 months)	0.393	0.009	1.343	0.023	0.376	0.411
Children with diarrhoea given ORS and HAF* (children age 0-59 months)	0.544	0.009	1.385	0.017	0.526	0.563
Households access to any improved source of drinking water	0.910	0.002	1.813	0.002	0.907	0.914
Households using improved sanitation facility	0.418	0.003	1.933	0.007	0.412	0.424
Households practicing open defecation	0.455	0.003	1.852	0.006	0.450	0.461
*Oral Re-hydration Solution (ORS) including Home Available Fluids (HAF)						